

1 **A.23 RIVER LAMPREY (*LAMPETRA***
 2 ***AYRESII*)**

3 **A.23.1 Legal and Other Status**

4 The river lamprey is a Class 3 (Watch List) California
 5 Species of Special Concern (Moyle et al. 1995, DFG 2011).
 6 It has no federal status.

7 **A.23.2 Species Distribution and Status**


8 **A.23.2.1 Range and Status**




photo courtesy Bradspictures.com

9 The river lamprey is known to occur from near Juneau, Alaska, to San Francisco Bay, California.
 10 The species appears to be more abundant in the lower Sacramento-San Joaquin River system
 11 than in other streams in California, but few surveys for river lamprey have been conducted
 12 (Moyle 2002). Population trends are unknown in California, although declines may have
 13 occurred due to freshwater habitat degradation. River lamprey are common in British Columbia,
 14 the center of their geographic range.

15 **A.23.2.2 Distribution and Status in the Plan Area**



16 California Department of Fish and Game has no records of river lamprey within Butte County (CNDDDB 2006); this species is expected to occur in the Plan Area (see Figure A-23). River lamprey are found in the Feather River from April through July upstream to the Fish Barrier Dam (SWRI 2003). This species was not reported in Butte Creek or Big Chico creeks or their tributaries, although the Pacific lamprey (*Entosphenus tridentata*) was found in Rock Creek, a tributary to Big Chico Creek (CSU Chico 1998, Big Chico Creek Watershed Alliance 1999). The species is also known to be present in the Sacramento River along the western boundary of the Plan Area (Information Center for the Environment no date).



1

2

3

Figure A-23. River Lamprey Modeled Habitat

1 **A.23.3 Habitat Requirements and Special Considerations**

2 The habitat requirements of river lamprey have not been well-studied. It is thought that adults need
3 clean, gravelly riffles in permanent streams to spawn successfully. The ammocoete stage requires
4 high quality, perennial backwaters or stream edges over a sandy substrate, into which they burrow,
5 with water temperatures below 25°C (Moyle et al. 1995). Although lamprey can pass barriers that
6 other fish cannot, large dams and other habitat modifications remain barriers to migration.

7 **A.23.4 Life History**

8 The biology of the river lamprey has not been studied in California; therefore, the following
9 section is based on the biology of river lamprey from British Columbia, which may not be the
10 same.

11 The river lamprey is anadromous and migrates from the ocean up rivers and streams to spawning
12 grounds. Adults enter freshwater in the fall and move upstream to suitable spawning areas with
13 perennial water (Moyle et al. 1995, Moyle 2002). They reach sexual maturity in streams, at
14 which time they may shrink in length by up to 20 percent. Spawning occurs from February
15 through May in gravelly riffles where the adults dig saucer-shaped depressions. Fecundity is not
16 well documented, but a study of two females in Cache Creek reported that one female produced
17 11,400 eggs and the other produced 37,300 eggs (Vladykov and Follett 1958). Adults die after
18 spawning. The eggs hatch into ammocoetes that remain in fresh water for approximately 3 to 5
19 years along sandy backwaters or stream edges where they bury in the sediments and feed on
20 algae and microorganisms. Water temperatures of less than 25°C are also required (Moyle et al.
21 1995). The ammocoetes begin to transform into adults during the summer at approximately 12
22 centimeters (cm) total length (TL). This process takes 9 to 10 months, and the new adults enter
23 the ocean in late spring. They spend approximately 3 to 4 months in the ocean where they grow
24 rapidly to 25-31 cm TL. River lamprey feed on a variety of host fish species in the 10-31 cm TL
25 range, but the most common prey appear to be herring and salmon. They feed by attaching to
26 their prey's back above the lateral line and eating the muscle tissue, even after the host fish dies
27 (Moyle 2002).

28 **A.23.5 Threats**

29 The primary threats to river lamprey are loss or degradation of habitat through dams, diversions,
30 pollution, stream channelization, and urbanization (Moyle et al. 1995).

31 **A.23.6 Relevant Conservation Efforts**

32 There have been few efforts to conserve river lamprey in the Central Valley of California. The
33 CALFED Ecosystem Restoration Program (ERP) designated the entire lamprey family as
34 “Enhance and/or Conserve” (CALFED Bay-Delta Program 2000). This designation indicates

1 that the ERP will undertake actions to conserve and enhance their abundance and distribution
2 and the community diversity in which they live for their long-term stability.

3 **A.23.7 Species Habitat Suitability Model**

4 **Juvenile Rearing Habitat/Migration/Spawning Habitat.** River lamprey juvenile rearing,
5 migration, and spawning habitat is defined as juvenile rearing, migration, and spawning habitat
6 delineated by DFG (CNDDDB 2007), SWRI (2003), and Information Center for the Environment
7 (no date). River lampreys use the mainstem Sacramento River along the western edge of the
8 Plan Area and in the Feather River as far upstream as the Fish Barrier Dam (SWRI 2003).

9 **Assumptions.** DFG and NMFS are the state and federal agencies responsible for managing river
10 lamprey and as such are considered to be the authority on the distribution of the species and its
11 habitat. Data gaps in DFG and NMFS information were augmented with information from
12 SWRI (2003), and Information Center for the Environment (no date). Although not well-studied,
13 it is thought that adult river lampreys need clean, gravelly riffles in permanent streams to spawn
14 successfully. The ammocoete stage requires high quality, perennial backwaters or stream edges
15 over a sandy substrate, into which they burrow, with water temperatures below 25°C (Moyle et
16 al. 1995). The scale of this habitat model precludes the ability to distinguish among the river
17 lamprey habitat types.

18 **A.23.8 Recovery Plan Goals**

19 A recovery plan has not been prepared for river lamprey because it is not federally listed as
20 threatened or endangered.

21 **A.23.9 References**

22 **Literature Cited**

23 Big Chico Creek Watershed Alliance. 1999. Existing Conditions Report. Accessed 14 December
24 2006. <http://www.bigchicocreek.org/nodes/aboutwatershed/ecr>.

25 CALFED Bay-Delta Program. 2000. Ecosystem Restoration Program Plan. Volume II:
26 Ecological Management Zone Visions. Final Programmatic EIS/EIR Technical
27 Appendix. Available at:
28 http://www.delta.dfg.ca.gov/erp/docs/reports_docs/ERPP_Vol_2.pdf.

29 CNDDDB (California Natural Diversity Database). 2007. California Department of Fish and
30 Game, Sacramento, CA.

31 CNDDDB (California Natural Diversity Database). 2006. Sacramento: California Department of
32 Fish and Game.

- 1 CSU Chico (California State University Chico). 1998. Butte Creek Watershed Project Existing
2 Conditions Report. Prepared for Butte Creek Watershed Conservancy.
- 3 DFG (California Department of Fish and Game). 2003. Species of Special Concern. Available at
4 www.dfg.ca.gov/hcpb/species/ssc/sscfish/sscfish.shtml. Revised 2 May 2003.
- 5 Information Center for the Environment. No date. Distribution Maps of Fishes in California.
6 <http://ice.ucdavis.edu/aquadiv/fishcovs/fishmaps.html>.
- 7 Moyle, P. B. 2002. Inland Fishes of California. Berkeley: University of California Press.
- 8 Moyle, P. B., R. M. Yoshiyama, J. E. Williams, and E. D. Wikramanayake. 1995. Fish Species
9 of Special Concern in California. 2nd ed. Prepared for California Department of Fish and
10 Game, Rancho Cordova. Contract No. 2128IF.
- 11 SWRI (Southwest Research Institute). 2003. Fish Distribution in the Feather River below
12 Thermalito Diversion Dam to the Confluence with the Sacramento River. Draft Report
13 SP-F3.2, Task 1 and SP-F21, Task 2. Oroville FERC Relicensing (Project No. 2100).
- 14 Vladykov, V. D. and W. I. Follett. 1958. Redescription of *Lampetra ayersi* (Gunther) of western
15 North America, a species of lamprey (Petromyzontidae) distinct from *Lampetra fluviatilis*
16 (Linnaeus) of Europe. *Journal of the Fisheries Research Board of Canada* 15(1):47-77.

17
18

1

This page intentionally left blank.

DRAFT